

# Golgi Body Encyclopedia Article

## Golgi Body

The following sections of this BookRags Literature Study Guide is offprint from Gale's For Students Series: Presenting Analysis, Context, and Criticism on Commonly Studied Works: Introduction, Author Biography, Plot Summary, Characters, Themes, Style, Historical Context, Critical Overview, Criticism and Critical Essays, Media Adaptations, Topics for Further Study, Compare & Contrast, What Do I Read Next?, For Further Study, and Sources.

(c)1998-2002; (c)2002 by Gale. Gale is an imprint of The Gale Group, Inc., a division of Thomson Learning, Inc. Gale and Design and Thomson Learning are trademarks used herein under license.

The following sections, if they exist, are offprint from Beacham's Encyclopedia of Popular Fiction: "Social Concerns", "Thematic Overview", "Techniques", "Literary Precedents", "Key Questions", "Related Titles", "Adaptations", "Related Web Sites". (c)1994-2005, by Walton Beacham.

The following sections, if they exist, are offprint from Beacham's Guide to Literature for Young Adults: "About the Author", "Overview", "Setting", "Literary Qualities", "Social Sensitivity", "Topics for Discussion", "Ideas for Reports and Papers". (c)1994-2005, by Walton Beacham.

All other sections in this Literature Study Guide are owned and copyrighted by BookRags, Inc.

# Contents

<a href="#">Golgi Body Encyclopedia Article.....</a>	<a href="#">1</a>
<a href="#">Contents.....</a>	<a href="#">2</a>
<a href="#">Golgi Body.....</a>	<a href="#">3</a>

# Golgi Body

The Golgi body, or Golgi apparatus is a collection of flattened membrane sacks called cisternae that carry out the processing, packaging, and sorting of a variety of cellular products in higher plants and animals. This important cellular organelle was named in honor of **Camillo Golgi**, the Italian neuroanatomist who first described it in brain cells late in the nineteenth century. An individual Golgi apparatus is usually composed of four to eight cisternae, each a micron or less in diameter stacked on top of each other like pancakes. Many cisternal stacks interconnected by tubules and mobile transport vesicles make up a Golgi complex, which often is located near the **nucleus** in the center of the cell. In some animal cells, this complex can be huge, filling much of the cytoplasmic space. In some plant cells, on the other hand, many small, apparently independent Golgi apparatuses are distributed throughout the cell interior.

Each Golgi stack has a distinct orientation. The *cis* or entry face is the site at which transport vesicles bringing newly synthesized products from the endoplasmic reticulum dock with and add their contents to the Golgi cisternae. A complex network of anastomosing (connecting) membrane tubules attach to and cover the fenestrated cisternae on the *cis* face and serves as a docking site for transport vesicles. From the *cis* face a flow of vesicles carry transport and chaperone proteins back to the endoplasmic reticulum, while secretory products move on into the medial cisternae where further processing takes place. Finally, the products move to the *trans* or exit face where they undergo final processing, sorting, and packaging into vesicles that will carry them to the cell surface for secretion or to other cellular organelles for storage or use. Complex oligosaccharides are synthesized in the Golgi apparatus, and glycoproteins are assembled as materials move through the compartments of this organelle. A unique set of **enzymes** and chaperone proteins occur in each of the Golgi compartments to direct and carry out this complex set of reactions.