

Heterozygous and Homozygous Encyclopedia Article

Heterozygous and Homozygous

The following sections of this BookRags Literature Study Guide is offprint from Gale's For Students Series: Presenting Analysis, Context, and Criticism on Commonly Studied Works: Introduction, Author Biography, Plot Summary, Characters, Themes, Style, Historical Context, Critical Overview, Criticism and Critical Essays, Media Adaptations, Topics for Further Study, Compare & Contrast, What Do I Read Next?, For Further Study, and Sources.

(c)1998-2002; (c)2002 by Gale. Gale is an imprint of The Gale Group, Inc., a division of Thomson Learning, Inc. Gale and Design and Thomson Learning are trademarks used herein under license.

The following sections, if they exist, are offprint from Beacham's Encyclopedia of Popular Fiction: "Social Concerns", "Thematic Overview", "Techniques", "Literary Precedents", "Key Questions", "Related Titles", "Adaptations", "Related Web Sites". (c)1994-2005, by Walton Beacham.

The following sections, if they exist, are offprint from Beacham's Guide to Literature for Young Adults: "About the Author", "Overview", "Setting", "Literary Qualities", "Social Sensitivity", "Topics for Discussion", "Ideas for Reports and Papers". (c)1994-2005, by Walton Beacham.

All other sections in this Literature Study Guide are owned and copyrighted by BookRags, Inc.

Contents

Heterozygous and Homozygous Encyclopedia Article.....	1
Contents.....	2
Heterozygous and Homozygous.....	3

Heterozygous and Homozygous

The term heterozygous refers to the situation where a specific **gene** is represented by two differing forms or alleles within the **diploid nucleus**. The opposite case within the **cell** is considered homozygous. A **homozygote** has both copies of the gene present in the same form, or as identical alleles. The alleles are present in the same loci, the same physical location on the chromosomes.

A **heterozygote** is capable of producing gametes of two different forms, one for each allele present. This is as a result of the **homologous chromosomes** moving to opposite ends of the dividing cell during **meiosis**. Conversely, the homozygote is only capable of producing the one type of **gamete** because there is only one type of allele present.

Phenotypically, the heterozygote and homozygote may be indistinguishable from each other. This effect is due to the phenomenon of dominance and recessiveness. If one allele is dominant, it will mask the observable effects of the recessive allele. It will only become clear that there are two different alleles present when the offspring are observed and the genetic make-up of the gametes is inferred. The dominant allele present in the homozygous form is phenotypically indistinguishable from the heterozygote. This form is referred to as homozygous dominant. The homozygous form of the recessive allele (homozygous recessive) has the phenotype of the recessive allele and it can be easily inferred from observation if it is known that that particular characteristic is recessive. There are cases where dominance is incomplete and the heterozygote shows a blending of the two alleles.

A heterozygote need not be a diploid as described above; polyploid heterozygotes can also exist. The recognizing characteristic of a heterozygote is that it does not breed true, or in other words, it produces two (for a diploid organism) or more (for a polyploid individual) forms of gametes. Polyploid homozygotes can also occur and are recognized by their ability to produce only the one type of gamete.